

## Measurement of Haemolysin Activities in *Vibrio vulnificus*

Hyun-Jung Lee<sup>1\*</sup>, Jeong-A Kim<sup>2</sup>, Mi-Ae Lee<sup>2</sup>, Soon-Jung Park<sup>3</sup> and Kyu-Ho Lee<sup>3</sup>

<sup>1</sup>National Cancer Institute, National Institutes of Health, Bethesda, USA; <sup>2</sup>Department of Life Science and Interdisciplinary Program of Integrated Biotechnology, Sogang University, Seoul, South Korea; <sup>3</sup>Department of Environmental Medical Biology and Institute of Tropical Medicine, Yonsei University College of Medicine, Seoul, South Korea

\*For correspondence: [hyun-jung.lee@nih.gov](mailto:hyun-jung.lee@nih.gov)

**[Abstract]** VvhA produced by *Vibrio vulnificus* exhibits cytolytic activity to human cells including erythrocytes. Since haemolysis by VvhA may provide iron for bacterial growth and pathogenicity, we investigated the expression of VvhA to elucidate the regulatory roles of Fur, a major transcription factor controlling iron-homeostasis. Fur repressed the transcription of *vvhBA* operon via binding to the promoter region. However, haemolysin content and haemolytic activity were lowered in cell-free supernatant of *fur* mutant. This discrepancy between the levels of *vvhA* transcript and VvhA protein in *fur* mutant was caused by exoproteolytic activities of the elastase VvpE and another metalloprotease VvpM, which were also regulated by Fur. *vvpE* gene expression was repressed by Fur via binding to the Fur-box homologous region. Regulation of VvpM expression by Fur did not occur at the level of *vvpM* transcription. *In vitro* proteolysis assays showed that both proteases efficiently degraded VvhA. In addition, the extracellular levels of VvhA were higher in culture supernatants of *vvpE* or *vvpM* mutants than in the wild type. Thus this study demonstrates that Fur regulates haemolysin production at the transcription level of the *vvhBA* operon and at the post-translation level by regulating the expressions of two VvhA-degrading exoproteases, VvpE and VvpM.

This protocol can be applied to other *Vibrio* strains with haemolysin activities, such as *Vibrio parahaemolyticus* (*V. parahaemolyticus*) or other human pathogen strains with similar haemolysin activities.

### Materials and Reagents

1. Bacterial strains
  - a. Wild type (*Vibrio vulnificus* MO6-24/O, Clinical isolate: biosafety level 2) with pRK415 plasmid (vector control)
  - b. *fur* mutant with pRK415
  - c. *fur* mutant with pRK415-*fur*

2. 1% human red blood cells (RBCs) (from healthy person who is a volunteer and diluted with PBS buffer to make a 1% RBC solution)
3. Tetracycline (3 mg/ml) (Sigma-Aldrich)
4. PBS buffer (Sigma-Aldrich, catalog number: P5368)
5. 0.02% Triton X-100 (Triton™ X-100 for molecular biology) (Sigma-Aldrich, catalog number: T8787)
6. LB broth (see Recipes)


### **Equipment**

1. Shaking incubator
2. Centrifuge (14,000 rpm or 21,000 x g)
3. Spectrophotometry (540 and 600 nm)
4. 0.22 micron filter

### **Procedure**

1. Grow 5 ml of bacterial cells at 30 °C shaking incubator, in modified Luria–Bertani (LB) [addition of NaCl to LB at a final concentration of 2.5% (w/v): *Vibrio vulnificus* is a marine bacterium, so 2.5% NaCl makes cell happy to grow] medium supplemented with antibiotics (tetracycline 3 mg/ml).
2. Take bacterial samples and measure cell density by spectrophotometry at 600 nm.
3. Collect the supernatants of cultures by centrifugation (14,000 rpm, 5 min, RT) and filtration (use with syringe filter: pore size 0.22 µm).
4. Prepare cell-free supernatants from each culture and serial dilute them with PBS by 1/2, 1/4 and 1/8 (final volume: 1 ml).
5. Incubate 1% RBC solution with the equal volume of the diluted supernatants at 37 °C for 1 h (no agitation).
6. After 1-h incubation, centrifuge of each samples to remove unlysed RBCs (14,000 rpm, 5 min, RT).
7. To make the fully lysis of 1% RBC solution, use equal volume of 0.02% Triton X-100 at 37 °C for 1 h (no agitation).
8. After centrifuge of each sample, take the upper part of samples. Measure the lysed RBCs samples by spectrophotometry at 540 nm, as described by Shinoda *et al.*, 1985 (for the blank at 540 nm, you can use culture broth without bacteria cells).
9. Calculate Haemolytic activity (HU) (HU was expressed as the reciprocal of the dilution factor showing 50% haemolysis) (Shinoda *et al.*, 1985).

## Representative data



**Figure 1. Haemolytic activities are decreased by *fur* mutation or iron deprivation.**

Haemolytic activities in the supernatants of wild type (pRK415),  $\Delta fur$  strain (pRK415) and  $\Delta fur$  strain (pRK415-*fur*) cultures. *V. vulnificus* strains were grown in LBS supplemented with 3  $\mu$ g/ml tetracycline for 2 h, and each culture was treated for 4 h with 2,2'-dipyridyl at a concentration of 0.1 mM (+ chelator) or 0 mM (no chelator). OD600 of the cultures of wild type (pRK415) and  $\Delta fur$  strain (pRK415-*fur*) were 2.0–2.5, and OD600 of the  $\Delta fur$  (pRK415) cultures were 1.0–1.5. Serial dilutions of cell-free supernatants were added to 1% RBC solution, and lysed RBCs were measured by spectrophotometry. Activity was expressed as haemolytic units (HU), the reciprocal of the dilution factor showing 50% haemolysis (Lee *et al.*, 2013).

## Recipes

- LB broth
  - 10 g of Peptone
  - 10 g of Yeast extract
  - 5 g Sodium Chloride per one liter

## Acknowledgments

This protocol has been adapted and modified from Shinoda *et al.* (1985) and Lee *et al.* (2013). This work was supported by the Mid-Career Researcher Program through a National Research Foundation grant funded by the Ministry of Education, Science and Technology,

Korea (No. 2009-0092822 to K.-H.L.) and by the Marine and Extreme Genome Research Center program of the Ministry of Land, Transport and Maritime Affairs, Korea (to K.-H.L.).

## References

1. Lee, H. J., Kim, J. A., Lee, M. A., Park, S. J. and Lee, K. H. (2013). [Regulation of haemolysin \(VvhA\) production by ferric uptake regulator \(Fur\) in \*Vibrio vulnificus\*: repression of \*vvhA\* transcription by Fur and proteolysis of VvhA by Fur-repressive exoproteases.](#) *Mol Microbiol* 88(4): 813-826.
2. Shinoda, S., Miyoshi, S., Yamanaka, H. and Miyoshi-Nakahara, N. (1985). [Some properties of \*Vibrio vulnificus\* hemolysin.](#) *Microbiol Immunol* 29(7): 583-590.