

Assay to Evaluate BAL Fluid Regulation of Fibroblast α -SMA Expression

Jennifer L. Larson-Casey¹ and A. Brent Carter^{1, 2, *}

¹Department of Medicine, Division of Pulmonary, Allergy, and Critical Care Medicine, University of Alabama at Birmingham, Birmingham, AL, USA; ²Birmingham Veterans Administration Medical Center, Birmingham, AL, USA

*For correspondence: bcarter1@uab.edu

[Abstract] Because transforming growth factor- β (TGF- β 1) induces differentiation of fibroblasts to myofibroblasts, we developed a protocol to evaluate alveolar macrophage-derived TGF- β 1 regulation of lung fibroblast differentiation (Larson-Casey *et al.*, 2016). The protocol evaluates the ability of mouse bronchoalveolar lavage (BAL) fluid to alter fibroblast differentiation. Fibroblast differentiation was measured by the expression of α -smooth muscle actin (α -SMA).

[Background] Alveolar macrophages play an integral role in pulmonary fibrosis development by increasing the expression of TGF- β 1 (He *et al.*, 2011). Our prior data demonstrate that alveolar macrophages are a critical source of TGF- β 1 as mice harboring a conditional deletion of TGF- β 1 in macrophages were protected from pulmonary fibrosis (Larson-Casey *et al.*, 2016). The expression of α -SMA is a defining feature of myofibroblasts, and TGF- β 1 is a well-characterized pro-fibrotic mediator that induces transformation of fibroblasts to myofibroblasts both *in vitro* (Desmoulière *et al.*, 1993) and *in vivo* (Sime *et al.*, 1997). Prior studies exposed fibroblasts to recombinant TGF- β 1 to show its effect on differentiation and function (Horowitz *et al.*, 2007). Here we have developed a protocol for determining the ability of mouse BAL fluid to alter the differentiation of human lung fibroblasts to myofibroblasts, the cells that produce extracellular matrix proteins.

Materials and Reagents

1. 6-well cell culture plates (Corning, Costar[®], catalog number: 3516)
2. Normal human fibroblasts (IMR-90) (ATCC, catalog number: CCL-186)
3. DMEM
4. Fetal bovine serum (FBS) (Thermo Fisher Scientific, Gibco[™], catalog number: 26140095)
5. Penicillin-streptomycin (10,000 U/ml, 10,000 μ g/ml) (Thermo Fisher Scientific, Gibco[™], catalog number: 15140122)
6. Amphotericin B (Thermo Fisher Scientific, Gibco[™], catalog number: 15290018)
7. RPMI 1640 medium, no phenol red (Thermo Fisher Scientific, Gibco[™], catalog number: 11835030)
8. DPBS (Thermo Fisher Scientific, Gibco[™], catalog number: 14190144)
9. NP-40

10. Sodium chloride (NaCl)
11. Protease inhibitor tablets (Sigma-Aldrich, catalog number: 11836170001)
12. Phosphatase inhibitor (EMD Millipore, catalog number: 524625)
13. α -SMA antibody (American Research Products, catalog number: 03-61001)
14. β -actin antibody (Sigma-Aldrich, catalog number: A5441)
15. Tween 20
16. Fibroblast culture media (see Recipes)
17. Lysis buffer (see Recipes)

Equipment

1. Cell culture incubator, 37 °C, 5% CO₂: 95% air atmosphere (Thermo Fisher Scientific, Forma™, model: Direct Heat CO₂ Incubator)

Procedure

1. Seed fibroblasts at a density of 1×10^6 per well of a 6-well cell culture plate in a total volume of 1 ml. Allow cells to adhere to cell culture plate (2-6 h).
2. Harvest bronchoalveolar lavage (BAL) fluid from mice (Han and Ziegler, 2013). Mice can be treated with bleomycin (1.75 U/kg) or saline as a negative control. Determine the protein concentration of bronchoalveolar lavage (BAL) fluid. Using RPMI 1640 media, normalize all BAL samples to the same protein concentration in a total volume of 1 ml.

Notes:

- a. Typically, 0.5-1 ml of BAL fluid is used and RPMI is used to normalize the protein concentration to a final total volume of 1 ml.*
- b. Alternatively, this assay can be performed in vitro using conditioned media from macrophage cell lines or bone marrow derived macrophages. Harvest the conditioned media from macrophages post-transfection or post-treatment, using 1 million cells per 1 ml of media.*
3. Remove media from fibroblasts and rinse with 1.5 ml room temperature 1x PBS, add normalized BAL fluid to fibroblasts.
 - a. Incubate fibroblasts with BAL fluid for 24 h in a cell culture incubator, 37 °C, 5% CO₂: 95% air atmosphere.
 - b. Harvest fibroblasts by lysing cells in lysis buffer (see Recipes).
4. Determine α -SMA expression by immunoblot analysis. The expression of α -SMA protein was determined by Western blotting using 20 μ g of total cellular protein. After blocking in 5% milk, the membranes were probed with mouse α -SMA primary antibody using a 1:3,000 dilution in Tris buffered saline containing 0.1% Tween 20, the membranes were incubated with the anti-mouse secondary antibody (1:2,000) in antibody dilution buffer for 1 h.

Data analysis

Results can be shown as a representative immunoblot (Figure 1); also see Larson-Casey *et al.* (2016).




Figure 1. Immunoblot analysis of α -SMA in IMR-90 fibroblasts cultured in BAL fluid (BALF) from saline or bleomycin (Bleo) exposed WT mice

Recipes

1. Fibroblast culture media
DMEM
10% heat-inactivated FBS
100 U/ml, 100 μ g/ml penicillin-streptomycin
1.25 μ g/ml amphotericin B (fungizone)
2. Lysis buffer
1% NP-40
0.15 M NaCl
0.05 M Tris pH 7.4
1 protease tablet
Phosphatase inhibitor diluted 1:100

Acknowledgments

This work was supported by 2R01ES015981 & VA merit review BX001135.

References

1. Desmoulière, A., Geinoz, A., Gabbiani, F. and Gabbiani, G. (1993). [Transforming growth factor-beta 1 induces alpha-smooth muscle actin expression in granulation tissue myofibroblasts and in quiescent and growing cultured fibroblasts.](#) *J Cell Biol* 122(1): 103-111.
2. Han, H. and Ziegler, S. F. (2013). [Bronchoalveolar lavage and lung tissue digestion.](#) *Bio-protocol* 3(16): e859.

3. He, C., Murthy, S., McCormick, M. L., Spitz, D. R., Ryan, A. J. and Carter, A. B. (2011). [Mitochondrial Cu,Zn-superoxide dismutase mediates pulmonary fibrosis by augmenting H₂O₂ generation](#). *J Biol Chem* 286(17): 15597-15607.
4. Horowitz, J. C., Rogers, D. S., Sharma, V., Vittal, R., White, E. S., Cui, Z. and Thannickal, V. J. (2007). [Combinatorial activation of FAK and AKT by transforming growth factor-beta1 confers an anoikis-resistant phenotype to myofibroblasts](#). *Cell Signal* 19(4): 761-771.
5. Larson-Casey, J. L., Deshane, J. S., Ryan, A. J., Thannickal, V. J. and Carter, A. B. (2016). [Macrophage Akt1 kinase-mediated mitophagy modulates apoptosis resistance and pulmonary fibrosis](#). *Immunity* 44(3): 582-596.
6. Sime, P. J., Xing, Z., Graham, F. L., Csaky, K. G. and Gauldie, J. (1997). [Adenovector-mediated gene transfer of active transforming growth factor-beta1 induces prolonged severe fibrosis in rat lung](#). *J Clin Invest* 100(4): 768-776.