

## Design and Direct Assembly of Synthesized Uracil-containing Non-clonal DNA Fragments into Vectors by USER™ Cloning

Morten Egevang Jørgensen<sup>1</sup>, Nikolai Wulff<sup>2, 3</sup>, Majse Nafisi<sup>3, 4</sup>, Deyang Xu<sup>2, 3</sup>, Cuiwei Wang<sup>2, 3</sup>, Sophie Konstanze Lambertz<sup>2, 3</sup>, Zeinu Mussa Belew<sup>2, 3</sup> and Hussam Hassan Nour-Eldin<sup>2, 3, \*</sup>

<sup>1</sup>University of Würzburg, Institute for Molecular Plant Physiology and Biophysics, Würzburg, Germany;

<sup>2</sup>DynaMo Center, Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Frederiksberg C, Denmark; <sup>3</sup>Copenhagen Plant Science Center, Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Frederiksberg C, Denmark; <sup>4</sup>Plant Biochemistry Laboratory, Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Frederiksberg C, Denmark

\*For correspondence: [huha@plen.ku.dk](mailto:huha@plen.ku.dk)

**[Abstract]** This protocol describes how to order and directly assemble uracil-containing non-clonal DNA fragments by uracil excision based cloning (USER cloning). The protocol was generated with the goal of making synthesized non-clonal DNA fragments directly compatible with USER™ cloning. The protocol is highly efficient and would be compatible with uracil-containing non-clonal DNA fragments obtained from any synthesizing company. The protocol drastically reduces time and handling between receiving the synthesized DNA fragments and transforming with vector and DNA fragment(s).

**Keywords:** USER™ cloning, Cloning of synthesized non-clonal DNA fragments, Fusion of DNA fragments, Uracil excision based cloning, uNCDFs, Geneart

**[Background]** For synthesized DNA, non-clonal linear DNA fragments (NCDF) have emerged as a cheaper and faster alternative to clonal fragments that are delivered sequenced and in a circular vector. NCDFs can be regarded as an IKEA solution to DNA synthesis where it is the customers that assemble their DNA fragments into a vector of choice and subsequently must verify the sequence of the final construct. In this protocol, we obtained uracil-containing NCDFs from Thermo Fisher Scientific Geneart, whose NCDFs are termed DNA strings. The uracil-containing NCDFs will here be named uNCDFs. In uNCDFs, uracils are inserted at designated positions during synthesis. We were able to clone uNCDFs directly into USER cloning compatible vectors by simply re-suspending the fragments in water, incubating them with linearized USER vector and USER enzyme (as described below) followed by transformation of *E. coli*. The procedure requires minimal handling, lasts less than 1 h from receipt to transformation and requires very small amounts of DNA. We have tested uNCDFs for almost a year in our laboratory and found them to perform consistently well. We recently cloned 13 transporter genes into a USER compatible *Xenopus* oocyte expression vector. Each gene fragment was obtained as a uNCDF (~2 kb in size). We tested a single colony for each cloning and found that 12 of the genes were inserted correctly into the pNB1u destination vector. For the last clone we found a single mutation which was correct in the second colony that we tested for that fragment (Jørgensen *et al.*, 2017). The pNB1u vector

is approximately 2.5 kb in size. In another project, we cloned a larger batch of uNCDFs composed of 64 fragments 1-2.4 kb in size into a variety of USER compatible vectors including the large (~10 kb) USER compatible pCambia based vectors (Nour-Eldin *et al.*, 2006). Of these, 55 were correct after testing a single colony, 62 were correct after sequencing a second colony, while the remaining two were found in a correct form in the third colony (unpublished). We also used the protocol to seamlessly fuse three fragments together and insert them into a destination vector by USER fusion (Geu-Flores *et al.*, 2007). Thus, we experienced high efficiency when cloning uNCDFs which was comparable to the efficiency of cloning uracil-containing DNA fragments generated via PCR. Moreover the uNCDFs were shown to work well for both small and large USER compatible vectors.

The innovation in this protocol is the ability to design and order synthesized DNA fragment containing uracils at appropriate locations. To appreciate the value of this advance and ensure an understanding of this cloning technique for researchers who have not used USER cloning previously, a brief introduction to USER cloning is given below.

The principle behind the efficiency of USER cloning lies in the ability to generate long, complementary overhangs. These overhangs can anneal to form stable hybridization products that can be used to transform *E. coli* without prior ligation. Most importantly, their generation is not dependent on the introduction of restriction sites. Hitherto, the generation of long single stranded overhangs has proceeded in two steps. First, a PCR reaction was performed on a DNA fragment of interest using primers containing a short 8-16 nt upstream extension that is preceded by a single deoxyuridine residue. The resulting PCR products are treated briefly with a commercial mix of uracil DNA glycosylase and DNA glycosylase Endo VIII. These enzymes, are included in the USER™ enzyme mix, and remove the two single deoxyuridine residues and enable the dissociation of the short, single-stranded fragments lying upstream from the cleavage sites. For the generation of overhangs in a USER compatible destination vector, a short cassette is inserted into it so that digestion with a restriction- and a nicking enzyme creates the desired long overhangs. The fact that long, custom-made overhangs can be generated on PCR products can be exploited to generate a series of PCR products with complementary overhangs. This enables the generation of a hybridization product consisting of a vector and multiple PCR products, which can be fused into a compatible vector as easily as a single PCR product. For more detailed information on the cloning method, readers are referred to the following references on how to perform USER cloning and USER fusion (Nour-Eldin *et al.*, 2006 and 2010; Geu-Flores *et al.*, 2007). Additionally, bioinformatic tools have been developed to aid design of primers and strategy of USER cloning and fusion including the (Automated DNA Modifications with USER cloning) AMUSER web server tool for automated primer design (Genee *et al.*, 2015) and an interactive lab simulation that teaches the principles of USER cloning (<https://www.labster.com/simulations/user/>). Finally, overlap design for USER fusion has recently been optimized (Cavaleiro *et al.*, 2015).

Introducing uracils into DNA fragments was hitherto performed via PCR using uracil-containing primers. The PCR was performed after receiving the NCDF and required designing and ordering of appropriate uracil-containing primers. Incorporating uracils in DNA fragments during synthesis omits the

need to design and order uracil-containing primers and also omits the need to perform a PCR reaction, this drastically reduces time and handling between receiving the synthesized DNA fragments and transforming with vector and DNA fragment(s).

## **Materials and Reagents**

### **A. For ordering uracil-containing NCDFs**

1. Sequence of DNA to be synthesized (optimized for desired organism)

*Note: Depending on the fidelity of synthesis, any size of fragment is in principle suitable for synthesis. Until now the maximum size of NCDFs we have been able to order is 3 kb (varies depending on the company) and the fidelity in terms of how many colonies we had to sequence to find a correct clone has generally been 1-2.*

2. Sequence of USER tails to be added for insertion into USER vector (for single gene insertion)
3. Sequence of USER overlaps for seamless USER fusion

*Note: For insertion into USER compatible vectors, the USER tails are given by the USER cassette that has been inserted. Please consult the relevant publications. For USER fusion we typically use USER overlaps between 7-16 bp in length.*

### **B. For linearizing USER compatible vector**

1. 1.5 ml centrifuge tubes (Eppendorf tubes)
2. USER compatible vector (Nour-Eldin *et al.*, 2006 and 2010; Geu-Flores *et al.*, 2007)
3. *PacI* restriction enzyme (New England Biolabs, catalog number: R0547S)
4. *Nt.BbvCI* nicking enzyme (New England Biolabs, catalog number: R0632S)
5. Cutsmart buffer (included when ordering enzymes above)
6. H<sub>2</sub>O
7. PCR purification kit of choice (e.g., QIAquick PCR purification kit) (QIAGEN, catalog number: 28104)

*Note: For part B, follow protocol published previously (Nour-Eldin *et al.*, 2010).*

### **C. For cloning uNCDFs into linearized USER compatible vector**

1. 1.5 ml centrifuge tubes (Eppendorf tubes)
2. CaCl<sub>2</sub> competent *E. coli* cells (DH5α or NEB10β or other common cloning strains)

*Note: Do not perform transformation by electroshocking as the shock will cause the vector and uNCDF to dissociate.*

3. uNCDFs from synthesis company (in this case from Thermo Fisher Scientific Geneart)  
*Note: Despite their success, uNCDFs are not yet offered officially by Geneart. Please contact e.g., Anja Martinez at ThermoFisher Geneart ([anja.martinez@thermofisher.com](mailto:anja.martinez@thermofisher.com)) or us for ordering questions.*

4. H<sub>2</sub>O

5. Linearized USER compatible vector

*Note: A long list of vectors has been made USER cloning compatible for almost all organisms. Search within these three papers (Nour-Eldin *et al.*, 2006 and 2010; Geu-Flores *et al.*, 2007) and the many papers citing them. Please note, we are currently in the process of gathering USER vectors from groups from all over the world in order to deposit them at Addgene for easy accessibility.*

6. USER enzyme (New England Biolabs, catalog number: M5505S)

7. 5x or 10x PCR buffer (any kind)

## **Equipment**

1. 37 °C heat block
2. 42 °C heat block/water bath

## **Procedure**

A. Design and order uracil-containing NCDFs for single fragment USER cloning

For cloning single uNCDFs into USER compatible vectors that have been generated by our lab, our generic 5' and 3' user tails are added to termini of the fragment to be inserted (5' USER tail: 5'-GGCTTAAU, 3' USER tail: 5'-GGTTTAAU) (for inserting into other USER compatible vectors please use the corresponding USER tails). When ordering synthesized DNA we typically codon optimize the sequence for the organism wherein the gene will be expressed. In an example below, we show how we ordered the gene AT1G15210 as a uNCDF from Thermo Fisher Scientific Geneart. As instructed by Geneart, we provided them with a word file containing sequences to be synthesized in FASTA format. At the beginning of the word document, we included general instructions to the synthesis company, which are given in italics below

*Dear synthesis company:*


*Green marks a T in the top strand that must be replaced by a U. Turquoise marks an A, which is complementary to a T on the opposite strand that must be replaced by a U (i.e., it is the T on the opposite strand that must be replaced by a uracil). Yellow marks sequence that may not be altered. Non-marked sequence represents coding sequence, which may be changed to overcome synthesis challenges. The gene will be expressed in *Xenopus laevis* oocytes. Please use the appropriate codon preference table for codon optimization and alternations.*

>AT1G15210

5'-GGCTTAATATGGACTACGATCC.....AACTTCCAGAGCCGCTAGATTAAACC-3'

*Note: Punctuation marks approx. 2 kb of sequence that was included in the order but which has been omitted here for brevity. In this example, yellow sequence represents our standard 5' and 3' USER tails.*

B. Cloning single uNCDs into linearized USER compatible vector



**Figure 1. Overview of the USER cloning technique.** A USER cassette in a USER-compatible vector (upper left corner) with a restriction site (Pacl, light blue) in the middle, one variable nt surrounding it (different for each side, yellow and green), and oppositely oriented nicking sites (Nt.BbvCl, tan). The USER vector is digested with Pacl and Nt.BbvCl to generate 8 nt overhangs. A DNA fragment (upper right corner) with uracils at appropriate positions can be generated by PCR as described previously or via synthesis as a uNCD as described here. Fragment and vector are mixed with USER<sup>TM</sup> enzyme mix (excising deoxyuridines, pink) and the digested USER-compatible vector. Following brief incubation, the hybridized product can be used to transform *E. coli* without prior ligation. This figure has been reproduced with permission from (Nour-Eldin *et al.*, 2006).

1. Each fragment is to be surrounded by USER tails that enable insertion into a USER compatible vector. In this example, we use our *X. laevis* expression vector pNB1u (Nour-Eldin *et al.*, 2006).

uNCDFs are mixed directly with the digested pNB1u vector without prior PCR amplification (Figure1).

- a. Dilute each uNCDF to 100 ng/μl in H<sub>2</sub>O (or TE buffer pH 8).
- b. The USER-compatible pNB1u *X. laevis* oocyte expression vector is digested with *PacI*/Nt.*BbvCI* overnight, PCR purified and diluted to a concentration of ~50 ng/μl (as previously described [Nour-Eldin *et al.*, 2006 and 2010]).

*Note: If a USER compatible vector is not available, it is possible to generate them via PCR. In that case, the vector backbone is treated as a DNA fragment to be assembled with DNA fragment of interest via USER fusion (please see below).*

- c. For the USER reaction, mix 100 ng uNCDF with 50 ng digested pNB1u, 1 U USER enzyme (NEB), 2 μl 5x PCR reaction buffer and 5 μl H<sub>2</sub>O.
- d. Incubate the reaction at 37 °C for 25 min, and then for 25 min at room temperature.
- e. Transform 50 μl chemically competent *E. coli* cells with the reaction mixture by heat shock (5 min on ice, 30-45 sec at 42 °C and 5 min on ice).
- f. Plate the transformation mixture on LB-plates containing appropriate antibiotic selection (for pNB1u we use carbenicillin or ampicillin-containing LB plates).
- g. Select three colonies from the plates and grow overnight. Extract plasmids and analyze the extracted plasmids by gel-electrophoresis. Sequence the plasmids with an insert.

#### C. Design and order uracil-containing NCDFs for USER fusion of multiple fragments

For fusion of multiple uNCDFs into USER compatible vectors overlap regions have to be selected and included in adjacent fragments. Overlap regions are selected as previously described by finding a T on the bottom strand and a T on the top strand 7-15 bases downstream of the first T. The pair of selected Ts should be within 20-30 bases distance to the junction site (Figure 2) (Geu-Flores *et al.*, 2007; Nour-Eldin *et al.*, 2010). These Ts will be the ones replaced by a U during synthesis. For insertion into the USER compatible vector, our generic 5' and 3' user tails are added to the terminal fragments. When ordering synthesized DNA we typically codon optimize the sequence for the organism wherein the gene will be expressed. In an example below, we show how we designed and ordered three fragments to be fused and inserted into a USER compatible vector by USER cloning. The full-length sequence (~7,200 bp) was split into three fragments (FAS1\_A, FAS1\_B and FAS1\_C). The sequence of their termini is given below. In each fragment, the punctuation in the middle denotes approximately 2 kb of sequence, which was included in the order but has been omitted here for brevity. Each fragment was designed to include overlap regions to the adjacent fragment close to the junction sites. Please see below and see (Geu-Flores *et al.*, 2007; Nour-Eldin *et al.*, 2010) for more detailed information on how to design USER fusion overlap regions. Alternatively, the Amuser web server tool can design overlap regions for any USER fusion approach (Genee *et al.*, 2015). As instructed by Geneart we provided them with a Word file containing sequences to be synthesized in FASTA format. At the beginning of the Word document, we included general instructions to the synthesis company, which are given in italics below.

Dear synthesis company:

Green marks a *T* in the top strand that must be replaced by a *U*. Turquoise marks an *A*, which is complementary to a *T* on the opposite strand that must be replaced by a *U* (i.e., it is the *T* on the opposite strand that must be replaced by a uracil). No alterations are allowed in any color shaded sequence. Non-marked sequence represents coding sequence, which may be changed to overcome synthesis challenges. The genes will be expressed in *Xenopus laevis* oocytes. Please use the appropriate codon preference table for codon optimization and alternations.

>FAS1\_A

5'-GGCTTAAUATGCCTGCTCTG.....AGGCTATTCTTAAAGAGAGCTTT-3'

*Note: Only the sense strand is included in the ordering process but the result is a uracil-containing double stranded fragment. No alterations are allowed in any colored region during the codon optimization process. Yellow marks our standard 5' USER tail. Green marks the *T*, which must be synthesized as *U*. Turquoise marks the *A*, whose complimentary *T* must be synthesized as *U*. Grey marks the overlap which will be exposed as a single stranded fragment upon USER treatment and which will hybridize to the complimentary single stranded overhang on the adjacent fragment (in this case FAS1\_B). Once synthesized, the double stranded FAS1\_A uNCDF will look as follows:*

5'-GGCTTAAUATGCCTGCTCTG.....AGGCTATTCTTAAAGAGAGCTTT-3'

3'-CCGAATTATACGGACGAGAC.....TCCGATAAGAAUTCTCTCGAAA-5'

Upon treatment with USER enzyme the double stranded FAS1\_A uNCDF will loose sequences lying upstream of the uracils and look as follows:

5'- ATGCCTGCTCTG.....AGGCTATTCTTAAAGAGAGCTTT-3'

3'-CCGAATTATACGGACGAGAC.....TCCGATAAGAA -5'

>FAS1\_B

5'-AAGAGAGCTTUGGGTCCTGAGGCTAC.....GAGTACCTGAAGAAAACCTTCCCT-3'

As for FAS1\_A only the sense strand is included in the ordering process. Green marks the *T*, which must be synthesized as *U*. Turquoise marks the *A*, whose complimentary *T* must be synthesized as *U*. Grey marks the overlaps which will be exposed as single stranded overhangs upon USER treatment and which will hybridize to the complimentary single stranded overhangs on the adjacent fragments (in this case the left 'grey' will hybridize to the overhang generated on FAS1\_A, whereas the right 'grey' will hybridize to the overhang generated on FAS1\_C). Once synthesized, the double stranded FAS1\_B uNCDF will look as follows:

5'-AAGAGAGCTTUGGGTCTGAGGCTAC.....GAGTACCTGAAGAAAACCTTCCCT-3'  
3'-TTCTCTCGAAACCCAGGACTCCGATG.....CTCATGGACTTCTTUGGAAGGGA-5'

Upon treatment with USER enzyme the double stranded FAS1\_B uNCDF will loose sequences lying upstream of the uracils and look as follows:

5'- GGGTCTGAGGCTAC.....GAGTACCTGAAGAAAACCTTCCCT-3'  
3'- TTCTCTCGAAACCCAGGACTCCGATG.....CTCATGGACTTCTT -5'

>FAS1\_C

5'-ACCTTCCCTCAGCTGACCGACG.....GGCAACGCTTAGATTAAACC-3'


As for FAS1\_A only the sense strand is included in the ordering process. Green marks the T, which must be synthesized as U. Turquoise marks the A, whose complimentary T must be synthesized as U. Grey marks the overlap, which will be exposed as single stranded overhang upon USER treatment and which will hybridize to the complimentary single stranded overhangs on the adjacent fragment (in this case the rightmost 'grey' overlap region on FAS1\_B). Once synthesized, the double stranded FAS1\_B uNCDF will look as follows:

5'-ACCTTCCCTCUGCTGACCGACG.....GGCAACGCTTAGATTAAACC-3'  
3'-TGGAAGGGAGTCGACTGGCTGC.....CCGTTGCGAATCUAATTGG-5'

Upon treatment with USER enzyme the double stranded FAS1\_C uNCDF will loose sequences lying upstream of the uracils and look as follows:

5'- GCTGACCGACG.....GGCAACGCTTAGATTAAACC-3'  
3'- TGGAAGGGAGTCGACTGGCTGC.....CCGTTGCGAATC -5'

#### D. Fusion and clone of multiple uNCDFs into linearized USER compatible vector



**Figure 2. Overview of the USER fusion technique.** Fragments X1, X2, and X3 are to be fused together. Overlap regions are marked in grey. Uracils can be inserted via PCR or as described here during synthesis. The DNA fragments are mixed with a pre-digested USER-compatible vector and treated with the deoxyuridine-excising USER<sup>TM</sup> enzyme mix. This generated 3' overhangs that complement each other (indicated by arrows), while the outermost ones complemented the overhangs of the pre-digested vector. This design enables the formation of a stable circular hybridization product that can be transformed directly into *E. coli* without prior ligation. This figure has been reproduced with permission from (Geu-Flores *et al.*, 2007).

For fusing and cloning multiple uNCDFs into USER compatible vectors that have been generated by our lab, our generic 8 bp long 5' and 3' user tails are added to termini of the terminal fragments. At junction sites uracil-containing overlaps were included to allow seamless fusion upon mixing. When possible we strive to choose fusion overlaps of different lengths and composition to minimize mis-hybridization between fragments. In this example, we fuse and clone three fragments into our *X. laevis* expression vector pNB1u (Nour-Eldin *et al.*, 2006). Each fragment contained a uracil at the appropriate location in each USER tail. The uracil was incorporated during synthesis. Thus, uNCDFs are mixed in equimolar ratios directly with the digested pNB1u vector without prior PCR amplification with uracil-containing primers.

1. Dilute each uNCDFs to 50 ng/ $\mu$ l in H<sub>2</sub>O.
2. The USER-compatible pNB1u *X. laevis* oocyte expression vector is digested with *PacI*/*Nt.BbvCI* overnight, PCR purified and diluted to a concentration of ~50 ng/ $\mu$ l (as previously described (Nour-Eldin *et al.*, 2006 and 2010)).
3. For the USER reaction, mix 100 ng of each uNCDFs with 50 ng digested pNB1u, 1 U USER enzyme (NEB), 2  $\mu$ l 5x PCR reaction buffer and H<sub>2</sub>O to bring the total volume to 10  $\mu$ l.

4. Incubate the reaction at 37 °C for 25 min, and then for 25 min at room temperature.
5. Transform 50 µl chemically competent *E. coli* cells (for example NEB10β by heat shock) with the reaction mixture (5 min on ice, 30-45 sec at 42 °C and 5 min on ice).  
*Note: Typically the volume of added USER reaction mixture should not exceed 10% of the competent *E. coli* volume (i.e., max 5 µl reaction mixture to 50 µl *E. coli* volume, 10 µl reaction mixture to 100 µl *E. coli* volume etc. Exceeding this ration can reduce transformation efficiency.*
6. Plate the transformation mixture on LB-plates containing appropriate antibiotic selection (for pNB1u we used carbenicillin-containing LB plates).
7. Select eight colonies from the plates and grow overnight. Extract plasmids and analyze the extracted plasmids by gel-electrophoresis. Sequence the plasmids with an insert.

## Notes

If insertion does not succeed in the first trial, we encourage users—in addition to the USER cloning—to clone the uNCDFs into a blunt-ended cloning vector (such as pJET) for safekeeping and future work.

## Acknowledgments

MEJ is supported by a grant from the Danish Council for Independent Research: DFF–6108-00122. DX, AP, CW, DV, SKL and HNN were funded by DNRF99 grant from the Danish National Research Foundation. ZNB was funded by Innovationfund Denmark J.nr.: 76-2014-3 and NW was funded by Human Frontier Science Program RGY0075/2015. The authors declare no conflicts of or competing interest.

## References

1. Cavaleiro, A. M., Kim, S. H., Seppala, S., Nielsen, M. T. and Norholm, M. H. (2015). [Accurate DNA assembly and genome engineering with optimized uracil excision cloning](#). *ACS Synth Biol* 4(9): 1042-1046.
2. Genee, H. J., Bonde, M. T., Bagger, F. O., Jespersen, J. B., Sommer, M. O., Wernersson, R. and Olsen, L. R. (2015). [Software-supported USER cloning strategies for site-directed mutagenesis and DNA assembly](#). *ACS Synth Biol* 4(3): 342-349.
3. Geu-Flores, F., Nour-Eldin, H. H., Nielsen, M. T. and Halkier, B. A. (2007). [USER fusion: a rapid and efficient method for simultaneous fusion and cloning of multiple PCR products](#). *Nucleic Acids Res* 35(7): e55.
4. Jorgensen, M. E., Xu, D., Crocoll, C., Ramirez, D., Motawia, M. S., Olsen, C. E., Nour-Eldin, H. H. and Halkier, B. A. (2017). [Origin and evolution of transporter substrate specificity within the NPF family](#). *Elife* 6: e19466.

5. Nour-Eldin, H. H., Geu-Flores, F. and Halkier, B. A. (2010). [USER cloning and USER fusion: the ideal cloning techniques for small and big laboratories](#). *Methods Mol Biol* 643: 185-200.
6. Nour-Eldin, H. H., Hansen, B. G., Norholm, M. H., Jensen, J. K. and Halkier, B. A. (2006). [Advancing uracil-excision based cloning towards an ideal technique for cloning PCR fragments](#). *Nucleic Acids Res* 34(18): e122.