

HIV_{GKO}: A Tool to Assess HIV-1 Latency Reversal Agents in Human Primary CD4⁺ T Cells

Emilie Battivelli^{1, 2, 3, \$} and Eric Verdin^{1, 2, 3, \$, *}

¹Gladstone Institute of Virology and Immunology, Gladstone Institutes, San Francisco, CA, USA;

²Department of Medicine, University of California San Francisco, San Francisco, CA, USA; ³Buck Institute for Research on Aging, Novato, CA, USA; ^{\$}Present address: Buck Institute for Research on Aging, Novato, CA, USA

*For correspondence: EVardin@buckinstitute.org

[Abstract] While able to suppress HIV replication in HIV infected individuals, combination antiretroviral therapy (ART) fails to eliminate viral latent reservoir, which consists in integrated transcriptional silenced HIV provirus. So far, identification of latently-infected cells has relied on activating cells to induce expression of HIV proteins which can then be detected. Unfortunately, this activation significantly changed the cell phenotype. We developed a novel HIV reporter, named HIV_{GKO}, that allows the purification of latently-infected cells in absence of reactivation. Indeed, latent cells can be identified by expression of the EF1a-driven mKO2 and lack of expression of the LTR-driven csGFP. This protocol can be used to study the effectiveness of LRAs (Latency Reversal Agents) in reactivating latent HIV in primary cells.

Keywords: HIV-1 latency, Latency reversal, Latency reversal agents, Reservoirs, Human cells, Dual-fluorescence reporter, HIV_{GKO}, Flow cytometry

[Background] The new version of dual-labeled virus (HIV_{GKO}), contains a codon-switched eGFP (csGFP) under the control of the HIV-1 promoter in the 5' LTR and a distinct, unrelated fluorescent protein mKO2 under the control of the cellular elongation factor one alpha promoter (EF1 α). It is important to use unrelated fluorescent proteins in those reporters due to recombination issues when using fluorescent proteins genetically related. Productively infected cells are thus mostly csGFP⁺ mKO2⁺ (some might only be GFP⁺), while latently infected cells are csGFP⁻ mKO2⁺. Flow cytometers such as the sorter Arial II allows the purification of pure infected population (productive, latent and/or uninfected), while the analyzer LSRII allows for the assessment of the transcriptional activation of the HIV_{GKO} LTR within a short time frame post-infection.

Materials and Reagents

A. Production of HIV_{GKO} in HEK293T cells

1. 182 cm² tissue culture flask (VWR, catalog number: 10062-864)
2. Tips
 - 0.1-10 μ l (Fisher Scientific, FisherbrandTM, catalog number: 02-681-440)
 - 1-200 μ l (Fisher Scientific, FisherbrandTM, catalog number: 02-707-502)

101-1,000 μ l (Fisher Scientific, FisherbrandTM, catalog number: 02-707-509)

3. Pipettes

2 ml aspirating pipettes (VWR, catalog number: 414004-265)

5 ml (VWR, catalog number: 89130-896)

10 ml (VWR, catalog number: 89130-898)

25 ml (VWR, catalog number: 89130-900)

4. 15 ml conical tube (VWR, catalog number: 89039-666)

5. 1.5 ml Eppendorf tubes (Fisher Scientific, catalog number: 05-408-129)

6. 50 ml conical tube (VWR, catalog number: 89039-658)

7. UltraClear Centrifuge Tubes 25 x 89 mm (Beckman Coulter, catalog number: 344058)

8. 50 ml centrifuge tube filtration (VWR, catalog number: 89220-710)

9. HEK293T cells (ATCC, catalog number: CRL-3216)

10. Plasmids:

a. HIV_{GKO} (Battivelli *et al.*, 2018)

b. HIV dual-tropic envelope (pSVIII-92HT593.1) (NIH AIDS Reagent Program, catalog number: 3077)

11. DMEM (Corning, catalog number: 10-013-CVR)

12. RPMI (Corning, catalog number: 10-040-CVR)

13. Fetal bovine serum (FBS) (Gemini Bio-Products, BenchMarkTM, catalog number: 100-106)

14. 100x penicillin/streptomycin (Corning, catalog number: 30-002-CI)

15. 1x PBS (Corning, catalog number: 21-031-CVR)

16. Trypsin-EDTA (Corning, catalog number: 25-053-CI)

17. Cell culture water (Corning, catalog number: 25-055-CV)

18. Chloroquine diphosphate salt (Sigma-Aldrich, catalog number: C6628)

19. HEPES (Sigma-Aldrich, catalog number: H3375)

20. Potassium chloride (KCl) (Sigma-Aldrich, catalog number: P9541)

21. Dextrose (Fisher Scientific, catalog number: BP350-1)

22. Sodium chloride (NaCl) (Sigma-Aldrich, catalog number: S3014)

23. Sodium phosphate dibasic (Na₂HPO₄) (Fisher Scientific, catalog number: BP332-500)

24. Calcium chloride (CaCl₂) (Sigma-Aldrich, catalog number: C1016)

25. Nuclease-free H₂O (Thermo Fisher Scientific, InvitrogenTM, catalog number: AM9937)

26. FlaQ Assay reagents (Gesner *et al.*, 2014)

27. 25 mM chloroquine (see Recipes)

28. 2x HBSS buffer (see Recipes)

29. 2 M CaCl₂ (see Recipes)

B. Isolation of human primary CD4⁺ T cells

1. 182 cm² tissue culture flask (VWR, catalog number: 10062-864)

2. Tips

0.1-10 μ l (Fisher Scientific, FisherbrandTM, catalog number: 02-681-440)

1-200 μ l (Fisher Scientific, FisherbrandTM, catalog number: 02-707-502)

101-1,000 μ l (Fisher Scientific, FisherbrandTM, catalog number: 02-707-509)

3. Pipettes

2 ml aspirating pipettes (VWR, catalog number: 414004-265)

5 ml (VWR, catalog number: 89130-896)

10 ml (VWR, catalog number: 89130-898)

25 ml (VWR, catalog number: 89130-900)

4. 50 ml conical tube (VWR, catalog number: 89039-658)

5. Blood or LRC

6. RosetteSepTM Human CD4⁺ T cell enrichment cocktail (STEMCELL Technologies, catalog number: 15062)

7. Histopaque 1077 (Sigma-Aldrich, catalog number: 10771-500ML)

8. RPMI (Corning, catalog number: 10-040-CVR)

9. Fetal bovine serum (FBS) (Gemini Bio-Products, BenchMarkTM, catalog number: 100-106)

10. 100x penicillin/streptomycin (Corning, catalog number: 30-002-CI)

11. 1x PBS (Corning, catalog number: 21-031-CVR)

12. Ammonium chloride (NH₄Cl) (Sigma-Aldrich, catalog number: A9434-500G)

13. Potassium bicarbonate (KHCO₃) (Sigma-Aldrich, catalog number: 237205-100G)

14. Ethylenediaminetetraacetic acid disodium salt dihydrate (Na₂EDTA) (Sigma-Aldrich, catalog number: E6635-100G)

15. Recombinant human Interleukin-2 (R&D Systems, catalog number: 202-IL-010)

16. AKC lysis buffer (see Recipes)

C. Infection of human primary CD4⁺ T-cells with HIV_{GKO}

1. 182 cm² tissue culture flask (VWR, catalog number: 10062-864)

2. Tips

0.1-10 μ l (Fisher Scientific, FisherbrandTM, catalog number: 02-681-440)

1-200 μ l (Fisher Scientific, FisherbrandTM, catalog number: 02-707-502)

101-1,000 μ l (Fisher Scientific, FisherbrandTM, catalog number: 02-707-509)

3. Pipettes

2 ml aspirating pipettes (VWR, catalog number: 414004-265)

5 ml (VWR, catalog number: 89130-896)

10 ml (VWR, catalog number: 89130-898)

25 ml (VWR, catalog number: 89130-900)

4. 50 ml conical tube (VWR, catalog number: 89039-658)

5. 15 ml conical tube (VWR, catalog number: 89039-666)

6. 96-well plate V-bottom (Thermo Fisher Scientific, NuncTM, catalog number: 249570) and lids (Thermo Fisher Scientific, catalog number: 263339)

7. Pipetting reservoirs (VWR, catalog number: 89094-662)
8. Isolated CD4⁺ T cells
9. Viral stock
10. RPMI (Corning, catalog number: 10-040-CVR)
11. Fetal bovine serum (FBS) (Gemini Bio-Products, BenchMarkTM, catalog number: 100-106)
12. 100x penicillin/streptomycin (Corning, catalog number: 30-002-CI)
13. Recombinant human Interleukin-2 (R&D Systems, catalog number: 202-IL-010)
14. Dynabeads human T-activator CD3/CD28 (Thermo Fisher Scientific, catalog number: 111.32D)

D. Sorting cells

1. Tips
 - 0.1-10 µl (Fisher Scientific, FisherbrandTM, catalog number: 02-681-440)
 - 1-200 µl (Fisher Scientific, FisherbrandTM, catalog number: 02-707-502)
 - 101-1,000 µl (Fisher Scientific, FisherbrandTM, catalog number: 02-707-509)
2. Pipettes
 - 2 ml aspirating pipettes (VWR, catalog number: 414004-265)
 - 5 ml (VWR, catalog number: 89130-896)
 - 10 ml (VWR, catalog number: 89130-898)
 - 25 ml (VWR, catalog number: 89130-900)
3. 50 ml conical tube (VWR, catalog number: 89220-710)
4. 15 ml conical tube (VWR, catalog number: 89039-666)
5. 96-well plate V-bottom (Thermo Fisher Scientific, NuncTM, catalog number: 249570) and lids (Thermo Fisher Scientific, catalog number: 263339)
6. Falcon round-bottom 5 ml tubes with 35 µm cell strainer cap (Corning, catalog number: 352235)
7. RPMI (Corning, catalog number: 10-040-CVR)
8. Fetal bovine serum (FBS) (Gemini Bio-Products, BenchMarkTM, catalog number: 100-106)
9. 100x penicillin/streptomycin (Corning, catalog number: 30-002-CI)
10. Recombinant human Interleukin-2 (R&D Systems, catalog number: 202-IL-010)
11. 1x PBS (Corning, catalog number: 21-031-CVR)
12. Ethylenediaminetetraacetic acid disodium salt dihydrate (Na₂EDTA) (Sigma-Aldrich, catalog number: E6635-100G)
13. Sodium azide (NaN₃) (Sigma-Aldrich, catalog number: S2002)
14. FACS buffer (see Recipes)

E. Drug treatment

1. Tips
 - 0.1-10 µl (Fisher Scientific, FisherbrandTM, catalog number: 02-681-440)
 - 1-200 µl (Fisher Scientific, FisherbrandTM, catalog number: 02-707-502)
 - 101-1,000 µl (Fisher Scientific, FisherbrandTM, catalog number: 02-707-509)

2. RPMI (Corning, catalog number: 10-040-CVR)
3. Fetal bovine serum (FBS) (Gemini Bio-Products, BenchMarkTM, catalog number: 100-106)
4. 100x penicillin/streptomycin (Corning, catalog number: 30-002-CI)
5. Recombinant human Interleukin-2 (R&D Systems, catalog number: 202-IL-010)
6. LRAs

F. Analysis of LRAs efficacy by flow cytometry.

1. RPMI (Corning, catalog number: 10-040-CVR)
2. Fetal bovine serum (FBS) (Gemini Bio-Products, BenchMarkTM, catalog number: 100-106)
3. 100x penicillin/streptomycin (Corning, catalog number: 30-002-CI)
4. Recombinant human Interleukin-2 (R&D Systems, catalog number: 202-IL-010)
5. 1x PBS (Corning, catalog number: 21-031-CVR)
6. Ethylenediaminetetraacetic acid disodium salt dihydrate (Na₂EDTA) (Sigma-Aldrich, catalog number: E6635-100G)
7. Sodium azide (NaN₃) (Sigma-Aldrich, catalog number: S2002)
8. LIVE/DEAD Violet Fixable Dead Cell Stain Kit (Thermo Fisher Scientific, catalog number: L34963) (small kit)
9. FACS buffer (see Recipes)

Equipment

1. Pipetman
2. FlinnpipetteTM F2 multichannel pipettes
5-50 µl (Thermo Fisher Scientific, catalog number: 4662050)
30-300 µl (Thermo Fisher Scientific, catalog number: 4662070)
3. Pipette-Aids
4. Tabletop centrifuge for Eppendorf tubes (Eppendorf, model: 5415D)
5. Vortex (VWR, catalog number: 10153-838)
6. Tabletop centrifuge for 96-well plates, Eppendorf, 15 ml and 50 ml tubes; used for spin infection (Beckman Coulter, model: Allegra X-14R)
7. High Speed Ultracentrifuge (Beckman Coulter, model: Optima L-60) with SW 28 Ti Swinging-Bucket rotor (Beckman Coulter, model: SW 28 Ti)
8. 5% CO₂ tissue culture incubator, 37 °C (Thermo Fisher Scientific, model: FormaTM Steri-CultTM CO₂ Incubators, catalog number: 3307)
9. DynaMagTM-5 Magnet (Thermo Fisher Scientific, catalog number: 12303D)
10. FACS AriaII (BD Biosciences, San Jose, CA)
11. FACS LSRII (BD Biosciences, San Jose, CA)

12. Hemacytometer (Hausser Scientific, catalog number: 3200) or automated cell counter (ORLFO Technologies, catalog number: MXZ001) with cassettes type M (ORLFO Technologies, catalog number: MXC001)
13. Biosafety cabinet 'Level 2'
14. Ice bucket (VWR, catalog number: 10146-298)

Software

1. FlowJo 10 or other versions (Tree Star)

Procedure

A. Production of HIV_{GKO} viral particles in HEK293T cells

Note: HIV_{GKO} particles are produced after calcium phosphate transfection in HEK293T cells.

1. HEK293T cell culture
 - a. HEK293T cells are cultured in DMEM medium (supplemented with 10% FBS, 1% penicillin/streptomycin) in 182 cm² tissue culture flasks in 20 ml medium.
 - b. For maintenance of the HEK293T culture, when approaching confluence (~80%) cells, aspirate media, wash once with PBS, then trypsinize (0.05% trypsin) and plate cells after 1/12 dilution in complete DMEM, in 182 cm² tissue culture flasks. HEK293T cells are split every 3 days.
 - c. One day before transfection, plate approximately 4 x 10⁶ of HEK293T cells per 182 cm² tissue culture flasks in 18 ml of complete DMEM.

Note: After thawing frozen cell vials, HEK293T cells are cultured for at least one week before transfecting them for virus production. In order to maximize viral particle production, HEK293T cells are never kept more than 4 weeks in culture. HEK293T cells should never be grown to 100% confluency as they lose their ability to be transfected.

2. Calcium phosphate transfection

Note: Protocol below describes the transfection of cells plated in one 182 cm² tissue culture flask. To utilize the maximum space available in the ultracentrifuge, transfet about 15 flasks.

- a. Remove all medium from the flask.
- b. Add 17 ml of fresh DMEM medium (supplemented with 10% FBS, 1% penicillin/streptomycin) containing a final concentration of 25 µM of chloroquine (do not add on cells but, at the bottom of the flask—Figure 1), slowly swirl the flask to distribute the solution evenly and, incubate for 30-60 min before transfection.
- c. Prepare DNA mix in nuclease-free H₂O in a 15 ml conical tube (45 µg of HIV_{GKO} and 25 µg of HIV-1 dual-tropic envelope [pSVIII-92HT593.1]).
- d. Add nuclease-free H₂O to the DNA mixture for a final volume of 1.5 ml.
- e. Add 1.75 ml of 2x HBSS buffer.

- f. Vortex mixture at a medium-high setting while slowly adding (dropwise) 220 μ l of 2 M CaCl₂ to the diluted DNA, and keep vortexing for about 45 sec.
- g. Incubate at room temperature for 10-30 min before adding the prepared DNA solution to the bottom of the flask.
- h. Slowly swirl the flask to distribute the solution evenly.
- i. Culture for 6-8 h at 37 °C.
- j. Remove supernatant.
- k. Add 17 ml of DMEM (supplemented with 10% FBS, 1% penicillin/streptomycin) medium (add at the bottom of the flask, as shown in Figure 1, to avoid detaching cells).
- l. Incubate for 48 h at 37 °C to allow viral production.
- m. Collect the supernatant into a 50 ml Falcon tube.
- n. Centrifuge for 20 min at 800 \times g at RT or 4 °C.
- o. Filter supernatant through a 50 ml centrifuge tube filtration.
- p. Transfer filtered supernatant into UltraClear Centrifuge Tubes (To avoid the collapse of the tube, make sure it is filled with at least 34 ml—add media or PBS to viral supernatant if necessary to reach that volume).
- q. Spin viral supernatant in the ultracentrifuge for 2 h at 100,000 \times g at 4 °C.
- r. Pour out supernatant, dry as much as you can the inside of the tube and resuspend the pellet with 50 μ l of cold RPMI (supplemented with 10% FBS, 1% penicillin/streptomycin) medium or FBS, make aliquots in 1.5 ml Eppendorf tubes and freeze at -80 °C.

Notes:

- a. *When scaling up the viral production, you should be able to see a pellet. Also, scale up the volume used to resuspend the pellet. Do not make air bubbles when resuspending the pellet and, pipet up and down at least 50 times.*
- b. *Concentrated viral supernatant can also be titered and used fresh. However, even though the p24 content will not change, fresh versus frozen virus will give different infection outcomes, which will require different viral input to start with (described in Step C2).*
- s. Thaw one aliquot, make viral dilutions up to 10,000 to 1 million and, titer virus for p24 content using the FlaQ assay protocol (Gesner *et al.*, 2014).

Note: In addition of titrating p24 content of the virus with the FlaQ assay, I would recommend directly titering the infection rate of your viral stock on activated cells using 4 or 5 viral dilutions, before proceeding with big experiments. You want to avoid total infection greater than 15%.




Figure 1. Adding media to cells without detaching cells. Steps A2b and A2k require to add fresh media with and without chloroquine, respectively. First, aspirate media, and then add fresh media directly at the bottom of the flask to prevent detaching the cells. Once the media is in the bottom of the flask, slowly swirl the flask to distribute the solution evenly and place the flask back into the incubator.

B. Isolation of human primary CD4⁺ T cells

1. Human primary CD4⁺ T cells culture

- a. Human primary CD4⁺ T cells are cultured in RPMI medium (supplemented with 10% FBS, 1% penicillin/streptomycin) + 20 U/ml of IL-2 in tissue culture flasks or plates at a concentration of 5×10^6 /ml, in a CO₂ incubator at 37 °C.
- b. Half of the medium is replaced with fresh RPMI medium (supplemented with 10% FBS, 1% penicillin/streptomycin) + 20 U/ml of IL-2 every other day.

2. Isolation of human primary CD4⁺ T cells

Note: This protocol describes CD4⁺ T cells isolation using the RosetteSep Human CD4⁺ T Cell Enrichment Cocktail. Any other CD4⁺ T cells isolation kit can be used, but the protocol might slightly differ. Always follow the manufacturer's protocol.

- a. Order fresh blood or LRC in advance to have it delivered on the day of the experiment.

Note: One blood/LRC corresponds to one donor. Three donors should be tested at least, in two different experiments at least.

- b. Transfer blood into a 50 ml Falcon tube (if using LRC, cut both extremities of the chamber with clean scissors and let the blood drops into the tube).
- c. Add 1,800 µl of RosetteSep Cocktail to sample, incubate at RT for 20 min, and mix sample by swirling the tube every 5 min.
- d. In a new 50 ml Falcon tube, add 10 ml of Histopaque-1077.
- e. Dilute sample with equal volume of PBS containing 2% FBS.

- f. Slowly and carefully layer diluted sample on density gradient medium to minimize their mixing.
- g. Centrifuge sample for 20 min at 800 x g, RT, with **brake off**.
- h. CD4⁺ T cells are contained in the white ring, below the plasma phase (Figure 2). Pipet the ring and transfer it to a new 50 ml Falcon tube.
- i. Wash cells by filling up the tube with PBS containing 2% FBS.
- j. Centrifuge for 3 min at 800 x g and discard supernatant.
- k. If pellet appears red, resuspend pellet with 15 ml of **cold** AKC lysis buffer (see Recipes), incubate for 2 min at RT.
- l. Fill the tube up to 50 ml with PBS containing 2% fetal bovine serum, centrifuge for 3 min at 800 x g, and get rid of the supernatant.
- m. If the pellet is still red, repeat Steps B2k-B2l, if not, proceed to Step B2n.
- n. Resuspend pellet (CD4⁺ T cells) with warm, fresh RPMI medium (supplemented with 10% FBS, 1% penicillin/streptomycin) + 20 U/ml of IL-2.
- o. Count cells using an automated cell counter or a hemacytometer.
- p. Culture cells at a concentration of 5 x 10⁶/ml (250 million cells are cultured into 185 cm² [flat]).

Note: Isolated CD4⁺ T cells are mainly resting and can be kept in culture as such for several days. Medium can be changed every 4 to 5 days.




Figure 2. Isolation of human primary CD4⁺ cells by histopaque density gradient. A. Slowly layer the blood (containing the RosetteSep Cocktail and diluted to half with PBS + 2% FBS) on top of histopaque density gradient. Centrifuge for 20 min at 800 x g, RT, with brake off. B. CD4⁺ T cells are contained in the white ring, below the plasma phase, but above the histopaque and red cell phases. Carefully pipet out the ring.

C. Infection of human primary CD4⁺ T-cells with HIV_{GKO}

Note: This protocol describes the activation of 250 million CD4⁺ T cells using Dynabeads Human T-

Activator CD3/CD28. Any other human CD4⁺ T cells activators kit can be used, but the protocol might slightly differ. Always follow the manufacturer's protocol.

1. Activation of human primary CD4⁺ T cells
 - a. Vortex the Dynabeads Human T-Activator CD3/CD28 in the vial.
 - b. Transfer 3.125 ml of Dynabeads (1 bead for 2 cells, which is half of the manufacturer's protocol) to a 15 ml tube.
 - c. Place the tube on a magnet for 1 min and discard the supernatant.
 - d. Remove the tube from the magnet and resuspend the washed Dynabeads with 10 ml of the CD4⁺ T cells to activate (at a concentration of 5×10^6 /ml), and add those 10 ml to the rest of the CD4⁺ T cells to activate. Transfer the whole suspension to an appropriate vessel (5 millions of activated cells can be cultured in one well of a 24-well plate).
 - e. Culture in a CO₂ incubator at 37 °C for three days as such (you should see cells aggregates due to activation).

2. Infection of human primary CD4⁺ T cells with HIV_{GKO}

Note: The spin-infection of 1 million activated CD4⁺ T cells requires 30 µl of a 10,000 ng p24^{Gag}/ml viral dilution to reach a total infection rate of 9%-12% (to keep the ratios of latent versus productive infections consistent, avoid infection rates greater than 15%). However, the total infection rate (productive + latent infections) might slightly change according to fresh versus frozen stocks, viral stocks themselves and donors. Thus, in addition of titering the virus with the FlaQ assay, I would recommend directly titering your viral stock on activated cells using 4 or 5 viral dilutions, before proceeding with big experiments. Using frozen viral stocks requires the use of bigger amount of virus to reach 9%-12% infection rate.

- a. Three days post-activation, mix cells and beads and, transfer cells to 15 ml tubes.
- b. Place the tube on a magnet for 1 min and transfer the supernatant into a new 50 ml tube.
- c. Centrifuge cells for 3 min at RT at 800 x g.
- d. Discard supernatant and resuspend the cell pellet of 250 million cells with media containing the viral preparation at 10,000 ng p24^{Gag}/ml.
- e. Transfer resuspended cells to a pipetting reservoir and distribute 30 µl of the cells/virus suspension per one well of a 96 well V-bottom plate using a multi-channel pipetman.
- f. Centrifuge (= spin-infection) the plate for 2 h, at 800 x g, 32 °C.
- g. Add 100 µl of fresh RPMI medium (supplemented with 10% FBS, 1% penicillin/streptomycin) + 20 U/ml of IL-2 in each well and pool cells back together. Resuspend cells in a final volume of 100 ml of fresh RPMI medium (supplemented with 10% FBS, 1% penicillin/streptomycin) + 20 U/ml of IL-2 and, transfer into a 182 cm² flask.

Note: Viral solution stays in, but it is possible to wash it away after spin-infection.

- h. Culture as such for 4 to 5 days in a CO₂ incubator at 37 °C.
- i. Replace half of the medium by centrifuging down cells, with fresh RPMI medium (supplemented with 10% FBS, 1% penicillin/streptomycin) + 20 U/ml of IL-2 every other day.

D. Sorting cells

1. Five days post-infection, collect and transfer cells to 50 ml tubes.
2. Centrifuge cells for 3 min at RT at 800 x g.
3. Discard supernatant and resuspend cells in 2 ml of FACS buffer (see Recipes).
4. Pipette resuspended cells through a 5 ml 35 µm cell strainer capped tube and place on ice.
5. Prepare several 15 ml collection tubes with 2 ml of FBS and place on ice. Do not forget to label your tubes.
6. Then proceed directly to cell sorting using the flow cytometer FACS Ariall (PE channel for mKO2, and FITC channel for GFP) (see Figure 3 for gating strategy).

Note: The sort can take up to 10 h, thus sort cells at 4 °C. Use the 85 µm nozzle to prevent spontaneous reactivation of latently infected cells. When collecting populations into 15 ml tubes, you can only collect 2 different populations at once. Keep sorting the latent population at all time, and exchange tubes for uninfected and productively infected cells when you have reached 5 million cells per population. You should be able to collect 1.5-2.5 million latently infected cells depending on your infection rate.

7. Spin down sorted cells for 3 min at RT, 800 x g.
8. Resuspend cell populations with fresh RPMI medium (supplemented with 10% FBS, 1% penicillin/streptomycin) + 20 U/ml of IL-2 and, distribute equally into a 96-well V-bottom plate.
Note: To test 5 LRAs, sorted populations should be divided equally into 5 wells. Given that each well contains 200 µl, the pellet should be resuspended in 1 ml final of RPMI media. Note that a few hundred thousand cells/well are enough to assay LRAs activity.
9. Let cells rest overnight in a CO₂ incubator at 37 °C.

E. Drug treatment

1. After 24 h incubation, prepare 2x LRAs dilutions (dilute drugs with complete RPMI).
2. Remove 100 µl of medium from each well and add 100 µl of 2x concentrated LRAs dilutions.
3. Incubate for 24 h in a CO₂ incubator at 37 °C.

F. Analysis of LRAs efficacy by flow cytometry

1. Twenty-four hours later, remove all medium from wells, and wash with FACS buffer.
2. Spin down cells for 3 min at RT, 800 x g and discard supernatant.
3. Stain cells with live/dead violet marker (1/1,000 dilution of the marker in FACS buffer, 100 µl/well, incubate for 10-15 min on ice in the dark).
Note: Live/dead violet marker is perfect since it does not overlap with FITC and PE channels, and thus no compensation is needed.
4. Wash cells once with FACS buffer and, proceed directly to flow analysis (see Figure 4 for gating strategy).

Notes:

- a. *Avoid fixing samples with PFA since it decreases fluorescence intensity.*

b. For flow cytometry, Ariall sorter was used to run samples since fluorescence intensity is higher. However, other flow cytometers such as LSRII or Calibur are also suitable for these experiments as long as they have the right filters.

Data analysis

1. Sorting Cells (Figure 3)




Figure 3. Gating strategy to sort out uninfected, productively and latently infected cells.

A. Set the gate on live cells. Cell viability is monitored by forward (FSC-Area) and side scatter (SSC-Area) analysis. B and C. Gate successively on singlets FSC-Area vs. FSC-Width, and SSC-Area vs. SSC-Width. D. Set the gate on GFP/FITC-Area⁺ to sort productively infected cells, or on GFP/FITC-Area⁻ vs. mKO2/PE-Area⁺ to sort latently infected cells, or to GFP/FITC-Area⁻ vs. mKO2/PE-Area⁻ to sort uninfected cells. Run briefly each sorted sample when the sort is over to check purity (usually > 90%).

2. Analysis of LRAs efficacy by flow cytometry (Figure 4)




Figure 4. Gating strategy to analyze LRAs efficacy on the reversal of HIV-1 latency. A. Set the gate cells (avoid the left bottom corner where it is mainly debris) based on size (FSC-Area) and granularity (SSC-Area) analysis. B. Gate on singlets (FSC-Area vs. FSC-Height). C. Gate on live cells (Live) (Pacific Blue-Area vs. FSC-Height). D. Quantify the number of GFP⁺ cells which is the number of productively infected cells (GFP/FITC-Area⁺ vs. FSC-Area). Deduce the % of GFP⁺ cells quantified in the control sample to obtain the number of latently reactivated cells by the LRA tested. Each LRA is tested at least on 3 independent HIV_{GKO} infected C donors.

Notes

The HIV_{GKO} construct has a defective envelop and requires the addition of exogenous envelop while producing viral particles. We use an HIV CXCR4 tropism envelop to target human primary CD4⁺ T cells. It is possible to pseudotype that construct with other envelops such as VSV-G or HIV CCR5 tropism envelops to target other cell types (Cavrois *et al.*, 2006).

Recipes

1. 25 mM chloroquine
25 mM chloroquine in PBS
2. 2x HBSS
50 mM HEPES
10 mM KCl
12 mM dextrose
280 mM NaCl
1.5 mM Na₂HPO₄
Adjust pH to 7.1
Note: The pH is crucial!
3. 2 M CaCl₂
2 M CaCl₂ in nuclease-free H₂O
4. AKC Lysis buffer
850 ml H₂O
8.02 g (150 mM) of NH₄Cl
1 g (10 mM) of KHCO₃
37.2 mg (0.1 mM) of Na₂EDTA
Adjust pH to 7.2
Add H₂O to 1,000 ml, and store at 4 °C for months
5. FACS buffer
2% FBS
2 mM EDTA
0.1% Sodium Azide
PBS

Acknowledgments

E.B. was supported by a post-doctoral fellowship from UCSF CFAR and a CHRP fellowship. E.V. was supported by funds from NIH 1R01DA030216, 1DP1DA031126, NIH/NIAID R01Ai117864 NIH/NIDA/1R01DA041742-01, NIH/NIDCR/1R01DE026010-01, and 5-31532. We would also like to

thank Marielle Cavrois and Herk Kasler, respectively directors of the Gladstone and Buck Flow Cores. The Gladstone Flow Core was funded by NIH Grants P30AI027763 and S10 RR028962 and by the University of California, San Francisco-Gladstone Institute of Virology and Immunology Center for AIDS Research (CFAR). We thank the amfAR Institute for HIV Cure Research.

Competing interests

The authors declare that there is no conflict of interest or competing interest regarding the publication of this article.

References

1. Battivelli, E., Dahabieh, M. S., Abdel-Mohsen, M., Svensson, J. P., Tojal Da Silva, I., Cohn, L. B., Gramatica, A., Deeks, S., Greene, W. C., Pillai, S. K. and Verdin, E. (2018). [Distinct chromatin functional states correlate with HIV latency reactivation in infected primary CD4⁺ T cells](#). *Elife* 7: e34655.
2. Cavrois, M., Neidleman, J., Kreisberg, J. F., Fenard, D., Callebaut, C. and Greene, W. C. (2006). [Human immunodeficiency virus fusion to dendritic cells declines as cells mature](#). *J Virol* 80(4): 1992-1999.
3. Gesner, M., Maiti, M., Grant, R. and Cavrois, M. (2014). [Fluorescence-linked antigen quantification \(FLAQ\) assay for fast quantification of HIV-1 p24^{Gag}](#). *Bio-protocol* 4(24): e1366.