

Correction Notice: Two Different Methods of Quantification of Oxidized Nicotinamide Adenine Dinucleotide (NAD⁺) and Reduced Nicotinamide Adenine Dinucleotide (NADH) Intracellular Levels: Enzymatic Coupled Cycling Assay and Ultra-performance Liquid Chromatography (UPLC)-Mass Spectrometry

Karina S. Kanamori^{1, #, *}, Guilherme C. de Oliveira^{1, #}, Maria Auxiliadora-Martins², Renee A. Schoon³, Joel M. Reid³ and Eduardo N. Chini¹

¹Signal Transduction and Molecular Nutrition Laboratory, Kogod Aging Center, Department of Anesthesiology and Perioperative Medicine, Mayo Clinic College of Medicine, Rochester, USA

²Fundação de Apoio ao Ensino, Pesquisa e Assistência do Hospital das Clínicas da Faculdade de Medicina de Ribeirão Preto da Universidade de São Paulo (FAEPA), São Paulo, Brazil

³Oncology Research, Mayo Clinic College of Medicine, Rochester, USA

*For correspondence: mendes.karina@mayo.edu

#Contributed equally to this work

On September 05, 2018, we noticed and issued a correction notice regarding one typo in our published protocol (<https://bio-protocol.org/e2937>). It is located in Procedure B Step 1d. Instead of writing a dilution of 1:10 it is mistakenly written as 1:100. The typo was to add another “0” in that dilution of 1:10.

Unfortunately, at the time we were submitting the article for publication a mistake was made in regards to Maria Auxiliadora-Martins’ affiliation. Also, as Maria’s research was supported by FAPESP, I would like to amend an acknowledgement referring this grant as well. I kindly ask that an amendment is included as follows:

The correct affiliation for Maria Auxiliadora-Martins is **Fundação de Apoio ao Ensino, Pesquisa e Assistência do Hospital das Clínicas da Faculdade de Medicina de Ribeirão Preto da Universidade de São Paulo (FAEPA), São Paulo, Brazil.**

The updated acknowledgement is: This work was supported in part by grants from the Ted Nash Long Life Foundation, and the Glenn Foundation for Medical Research via the Paul F. Glenn Laboratories for the Biology of Aging at the Mayo

Clinic, a grant from Calico Laboratories, the Mayo Foundation, National Institutes of Health (NIH) grants from the National Institute of Aging (NIA, grant AG-26094), the Pancreatic Cancer SPORE project from NIH/NCI to E.N.C (grant CA102701-08), the Mayo Clinic Center for Cell Signaling in Gastroenterology (NIDDK P30DK084567). **Maria Auxiliadora-Martins was supported by a grant from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) Brazil (#2016/20089-4).**

References

1. Kanamori, K. S., de Oliveira, G. C., Auxiliadora-Martins, M., Schoon, R. A., Reid, J. M. and Chini, E. N. (2018). [Two different methods of quantification of oxidized nicotinamide adenine dinucleotide \(NAD⁺\) and reduced nicotinamide adenine dinucleotide \(NADH\) intracellular levels: Enzymatic coupled cycling assay and Ultra-Performance Liquid Chromatography \(UPLC\)-Mass Spectrometry.](#) *Bio-protocol* 8(14): e2937.