

Mouse Macrophage Differentiation by Induction with Macrophage Colony-Stimulating Factor

Ying Liu¹, Keqiang Chen¹, Chunyan Wang¹, Wanghua Gong², Teizo Yoshimura², Ji Ming Wang^{2*},
Mingyong Liu^{2*}

¹Laboratory of Molecular Immunoregulation, Cancer Inflammation Program, Center of Cancer Research, National Cancer Institute-Frederick, NIH, Frederick, USA; ²Basic Research Program, SAIC-Frederick, Frederick, USA

*For correspondence: liu_mingyong@yahoo.com.cn

[Abstract] Macrophages are differentiated from circulating blood monocytes and act as tissue-resident professional phagocytes. Macrophages function in both innate and adaptive immune systems of vertebrate animals. The cytokine macrophage colony-stimulating factor (M-CSF) is essential for the proliferation and differentiation of monocytes. Here, we described a simple method to induce the differentiation of mouse bone marrow-derived myeloid precursor cells into macrophages in the presence of M-CSF.

Materials and Reagents

1. Mice of interest
2. 1x DPBS (Lonza, catalog number: 17-512F)
3. ACK lysing buffer (Lonza, catalog number: 10-548E)
4. Dulbecco's Modified Eagle's Medium with high glucose and without L-glutamine (DMEM) (Lonza, catalog number: 12-614F)
5. Fetal Calf Serum (FCS) (Thermo Fisher Scientific, catalog number: SH30070.03)
6. Murine M-CSF (Peprotech, catalog number: 315-02)
7. 100x Penicillin/Streptomycin (Life Technologies, Invitrogen™, catalog number: 10378-016)
8. Complete DMEM medium (see Recipes)

Equipment

1. 40 µm nylon strainer (BD Biosciences, Falcon®, catalog number: 352340)
2. Needle (Gauge #25)
3. Centrifuge

Procedure

1. Mouse bone marrow (BM) cells are harvested from femurs by syringe and needle (Gauge #25) with 5 ml 1x DPBS (cut two ends of femurs and rinse out the cells by 1x DPBS).
2. Centrifuge the cell suspension at 200 x g at room temperature (RT) for 5 min.
3. Remove the supernatant and suspend the cell pellet with 2 ml ACK lysing buffer for 1 min to deplete red blood cells.
4. The cell suspension is directly filtered through a 40 µm nylon strainer (observe the cell lysate under the microscope to determine whether red blood cells are depleted completely. If there are still too many red blood cells, add more ACK lysing buffer until there are very few red blood cells).
5. Wash the strainer with 2 ml 1x DPBS and centrifuge the filtered cell suspension at 200 x g at RT for 5 min.
6. The cell pellet is washed once again with 1x DPBS and re-suspended in 15 ml complete DMEM medium with 20 ng/ml murine M-CSF in a 100 mm Petri dish (one femur cells per dish) in an incubator (37 °C, 5% CO₂).
7. After 3 days, half of the medium is replaced with fresh complete DMEM medium.
8. On Day 4, the cells containing more than 80% CD11b⁺/F4/80⁺ macrophages are ready for further characterization and functional experiments.

Recipes

1. Complete DMEM medium (500 ml)
 - 100 ml FCS
 - 5 ml 100x Penicillin/Streptomycin
 - 400 ml DMEM

Acknowledgments

This protocol has been adapted from Liu *et al.* (2013).

References

1. Liu, Y., Chen, K., Wang, C., Gong, W., Yoshimura, T., Liu, M. and Wang, J. M. (2013). [Cell surface receptor FPR2 promotes antitumor host defense by limiting M2 polarization of macrophages.](#) *Cancer Res* 73(2): 550-560.